

## REVIEW

## Central effects of the somatotrophic system

Harald Jörn Schneider, Uberto Pagotto<sup>1</sup> and Günter Karl Stalla*Max Planck Institute of Psychiatry, Department of Endocrinology, Kraepelinstr. 10, 80804 Munich, Germany and <sup>1</sup>Ospedale Sant'Orsola Malpighi, Operative Unit of Endocrinology, Department of Internal Medicine and Gastroenterology, Via Massarenti, 9, 40138 Bologna, Italy**(Correspondence should be addressed to G K Stalla; Email: stalla@mpipsykl.mpg.de)*

### Abstract

The somatotrophic axis interacts with the central nervous system (CNS) on several levels. Growth hormone (GH) and insulin-like growth factor-I (IGF-I) receptors are expressed in many brain areas including the hippocampus, pituitary and hypothalamus. GH and IGF-I can pass the blood-brain barrier by an as yet not completely understood mechanism. They can also be produced in the brain and thus act via paracrine/autocrine mechanisms. GH and IGF-I are important factors in the development and differentiation of the CNS and have protective properties in dementia, and in traumatic and ischemic injury of the CNS. An improvement in cognitive functioning in GH-deficient patients by GH substitution has been shown. Significant results could, however, only be achieved with supraphysiological doses. In some studies, a correlation between IGF-I and cognitive function in the elderly has been found. GH has an important impact on mood and well-being with GH secretory capacity being reduced in depression. Pulsatile GH secretion is closely related to slow wave sleep (SWS) with SWS being stimulated by GH releasing hormone and rapid eye movement (REM) sleep by GH.

*European Journal of Endocrinology 149 377–392*

### Introduction

The positive effects of growth hormone (GH) substitution on metabolism, the cardiovascular system and body composition have been described (1–6). Recently, the effects of the somatotrophic axis on the central nervous system (CNS) have come to be the focus of interest. However, it remains largely unclear whether these effects are mediated directly by GH or by its mediator – insulin-like growth factor-I (IGF-I) or, in some cases, by the hypothalamic neuropeptide growth hormone releasing hormone (GHRH). It is also unclear which functions are influenced by the above-mentioned hormones of the somatotrophic axis.

In humans, GH binding sites, indicative of GH receptors (GH-R) can be found in most areas of the CNS. They are detected in their highest concentrations in the choroid plexus, but are also found in relevant amounts in the hippocampus, putamen, thalamus, pituitary and hypothalamus (7–9). GH-R messenger ribonucleic acid (mRNA) has also been detected in samples of human fronto-parietal and temporal cortex (10). A reduction in GH binding sites in the brain is seen with increasing age (7, 8).

The question as to whether GH can pass the blood-brain barrier (BBB) is still a subject of discussion. In acromegalic patients with excessive GH levels in the plasma, some authors have found normal GH levels in the cerebrospinal fluid (CSF) (11), whereas others have found increased levels (12). When GH is applied

externally for GH substitution, there is an increase in GH levels in the CSF (13, 14). In rats, GH has been found in the cortex, hippocampus, thalamus and amygdaloid nucleus (15, 16). Active transport mechanisms, similar to the ones known for leptin and insulin, have been proposed (8, 17). It has been shown that a transport mechanism across the BBB exists for prolactin, whereby prolactin receptors in the choroid plexus act as transport molecules (18). GH binding sites are also found in highest concentration in the choroid plexus, so it is possible that they act as a transport mechanism in a similar fashion. In addition, GH mRNA can be found in the CNS, suggesting that GH also acts via auto-crine/paracrine mechanisms (19).

Similarly, IGF-I receptors are found predominantly in the hippocampus and parahippocampal areas, but also in the amygdala, cerebellum and cortex (20). There is evidence that IGF-I is transported across the BBB via transcytosis (17). In rat brain neurons, uptake of IGF-I could be shown (21). A relevant amount of IGF-I is produced in the brain with IGF-I mRNA being found predominantly in the brain stem and cerebellum in adult rats (22). GHRH receptors have been found in abundance in the cortex and brain stem of rats (23).

GH release occurs in a pulsatile fashion with most being secreted during the early hours of night sleep in males, but in a more variable pattern and with a higher output during the day in females (24). GH secretion underlies the regulation of GHRH and somatostatin, as well as ghrelin, the newly discovered


natural ligand of the GH secretagog receptor (GHS-R) (25) (see Fig. 1). GHRH stimulates (26) and somatostatin inhibits (27, 28) GH release. Ghrelin is a peptide hormone that is mainly produced in the stomach (25). The GHS-R is expressed in the hypothalamus (29) and in the pituitary (30), and ghrelin mRNA is found in the pituitary (30). Ghrelin is a potent stimulator of GH release (31–34) and acts synergistically with GHRH on GH release (34, 35). Additionally, it appears to be an important stimulus of appetite and food intake (36–39) that might be mediated by stimulation of agouti-related protein expression in the hypothalamus (29), and correlates negatively with body weight (40, 41). IGF-I is mainly produced in the liver and is stimulated by GH, but a certain amount is also directly produced in the target tissues. Additionally, it underlies the regulation of the feeding state. In anorexia IGF-I is reduced and correlates positively with body mass index (42) whereas in obesity a slight reduction in IGF-I and a negative correlation with body mass index has been found (43, 44). Central functions that

can be influenced by GH, IGF-I and GHRH include sleep, cognitive functions, mood, and neuroprotection. The aim of this review is to give an overview of the impact of the somatotrophic axis on these functions with a focus on clinical symptomatology.

## Neuroprotection

### Growth hormone

The somatotrophic axis plays a central role in the development and growth of the CNS. The distinction between GH- and IGF-I-mediated effects is often difficult; however, transgenic mouse models have shown that GH and IGF-I may exert distinct effects on the nervous system. For example, in transgenic mice overproduction of GH induces an increase in body size and motoneuron size. On the other hand, overproduction of IGF-I only induces an increase in body size, with the motoneuron size remaining unchanged (45).



**Figure 1** Modes of action on the brain and regulation of the somatotrophic system. GHRH stimulates GH release. GH release is also stimulated by the natural GH secretagog ghrelin synergistically with GHRH. GH stimulates IGF-I release in the liver that exerts a negative feedback on GHRH and GH. Both IGF-I and GH may pass the blood-brain barrier and exert distinct effects in the brain. Both hormones are also produced directly in the brain and might act via autocrine/paracrine mechanisms.

In spinal chord injuries a decreased production of GH and IGF-I can be found (46). In rats with spinal chord injuries the application of GH improved neurological functions and reversed the traumatic suppression of evoked potentials in the spinal chord (9). Scheepens *et al.* (47) studied rats with hypoxic brain damage: in areas with strong cell loss, they found pronounced immunoreactivity for GH. By intracerebroventricular application of GH the amount of neuronal loss could be reduced in the frontoparietal cortex, hippocampus and dorsolateral thalamus, but was left unchanged in the striatum. This matches the regional distribution of GH receptors and thus supports the hypothesis that these particular neuroprotective effects are mediated directly by GH, and not by IGF-I.

### **Insulin-like growth factor-I**

IGF-I supports the myelination of the CNS by stimulating proliferation and differentiation of oligodendrocytes. It is involved in the differentiation of neurons to specific cell types; it can increase levels of neurotransmitters, neurotransmitter receptors and proteins of the cytoskeleton; it can inhibit apoptosis in neurons (22) and it stimulates dendrite growth (48). Disruption of the IGF-I gene leading to loss of function induces neuronal loss in the hippocampus and striatum (49). In animal models it has been shown that IGF-I has a positive influence on markers of Alzheimer's disease such as cholinergic dysfunction, amyloid toxicity, tau-phosphorylation, and glucose metabolism (50).

In some regions of the mammalian brain, for instance in the dentate gyrus of the hippocampus, there is lifelong neurogenesis. With increasing age, the rate of neurogenesis decreases. This decrease is dependent on environmental factors, hormones, and growth factors such as IGF-I. It has been shown that in aged rats a 60% reduction in the differentiation of new cells to neurons occurs. This reduction could be reversed by the intracerebroventricular application of IGF-I (51). This is particularly interesting as IGF-I production and IGF-I receptor density also declines with increasing age. Thus, it can be speculated that the age-dependent decline of IGF-I and IGF-I receptors might be a possible factor contributing to the development of cognitive deficits seen in the elderly.

In experimental infarction by occlusion of the medial cerebral artery in rats, a 60% reduction in infarct volume and an improvement in neurological functions could be achieved by intranasal application of IGF-I (52, 53). Recently, the Torres-Aleman group has published some very interesting results. They studied the effects of treadmill running and IGF-I application in rats. They found that both exercise and IGF-I application elevate IGF-I levels in the cerebrospinal fluid and induce an increase in neuronal c-fos expression and hippocampal brain-derived neurotrophic factor (BDNF) (21), and exercise also induces an increase in neuron number in

the hippocampus (54). In addition, exercise prevents neuronal loss in toxic damage of the hippocampus and the brain stem and has the ability to reverse neurological deficits (55). All effects of exercise and IGF-I could be reversed by blocking IGF-I or the IGF-I receptor.

These results show that both IGF-I and exercise have neuroprotective abilities, that the neuroprotective potential of exercise is mediated by IGF-I, that IGF-I can pass the BBB under physiological conditions, and that c-fos and BDNF are possibly involved in the mediation of the neuroprotective effects of IGF-I.

These data show that GH and IGF-I are not only important factors for the growth and development of the CNS but also exert neuroprotective effects throughout life. However, the clinical relevance of these findings has scarcely been investigated and clinical studies in humans are advisable.

### **Cognitive functions**

The question of the influence of the somatotrophic system on cognitive functions has been the subject of debate for a long time. In healthy volunteers it could be shown that a single dose of GHRH improves memory function (56). Neuroprotective properties of GH and IGF-I, especially in the hippocampus, support the assumption that the somatotrophic axis has effects on cognitive abilities. An increase in aspartate, the ligand of the N-methyl-D-aspartate (NMDA) receptor has been found in the CSF of GH-deficient patients after GH replacement (57). NMDA receptor activation has been implicated both in memory function (58) and in attentional performance (59).

### **Growth hormone deficiency**

A higher incidence of school under-achievement in short-statured children, independent of etiology has been reported (60). However, it remains unclear whether this is due solely to psychosocial or physiological reasons, or is also dependent on GH deficiency (GHD). In 1996, Sartorio *et al.* published a review on the relationship of growth hormone and body size to cognitive abilities (60). Short children are often low achievers in reading, spelling and arithmetic, despite a normal IQ. Children with GHD show learning and attention deficits (60) and disturbance in visual-motor integration (61) in spite of a normal IQ. In a study comparing children with GHD to short children with normal GH secretion, no differences in a multifactorial intelligence test and several personality questionnaires could be found (62). This supports the assumption that other factors besides GHD play an important role in the occurrence of school under-achievement in short-statured children.

However, other studies underline the role of GH in cognitive function and life achievements. In adult patients with GHD there is an increased rate of unemployment

even though these patients show normal intelligence and scholastic achievements (60). An impaired social status in patients with childhood-onset (CO)-GHD has been reported compared with short (63) and normal controls (64). Impairment in memory and executive function has been found in patients surgically treated for pituitary tumors (65), and women with untreated GHD exhibited lower scores in neuropsychological tests than healthy controls (66).

Deijen *et al.* examined cognitive functioning in 31 men with multiple pituitary hormone deficiency (MPHD), 17 men with isolated CO-GHD and in a healthy control group (67). They found reduced cognitive abilities in many fields in MPHD patients compared with controls, but in patients with isolated GHD only an impaired memory function was found. This shows that GHD appears to have a direct influence on memory function whereas other cognitive impairments are due to additional hormone deficiencies.

### **Effects of growth hormone substitution**

In recent years some very interesting studies of GH substitution on cognitive function in GH-deficient subjects have been conducted (see Table 1). Deijen *et al.* studied the effects of three different doses of GH (1, 2, and 3 IU/m<sup>2</sup>/day) compared with a placebo treatment over six months, and afterwards in an open-label phase over 18 months. In the first phase, they found improvements in memory function (associate recognition task, associate learning task) compared with placebo only in the groups with high GH doses leading to supranormal IGF-I levels. In the open-label phase, after one and again after two years, they found improvements in memory functions at all doses (68). This study confirmed the positive findings of GH substitution on cognitive abilities in previously conducted studies with small patient numbers (69, 70).

In nine patients with GHD treated with GH (0.25 IU/kg/week) or placebo for 6 months and then with only GH for another 6 months, a significant improvement in measures of attention (digit backwards, verbal fluency, cognitive efficiency) could be shown (71). This is consistent with results from our institute. We conducted a double-blind placebo-controlled trial with 18 GHD patients (16 adulthood-onset (AO)-GHD, two CO-GHD, six women, 12 men). In the first phase, patients received either GH (0.25 IU/kg/week) or placebo for 6 months. In the second phase, all patients received GH for another 6 months. Patients treated with GH showed continuous improvement over 12 months in attention trials (digit cancellation test, trail-making test, see Fig. 2) but no differences in verbal short-term or long-term memory were noted (72).

Baum *et al.* studied adult patients with AO-GHD in a double-blind placebo-controlled GH replacement study (0.042–0.126 IU/kg/week, adjusted to IGF-I) over

18 months (73). Tests for intelligence, language, memory, abstract thinking and personality showed normal results. Mild impairments in verbal learning and visual memory were found but these were still in the normal range. In none of the measured parameters was there an effect of GH compared with placebo (73). In another trial by Degerblad *et al.* conducted in a cross-over design with six patients over 12 weeks, no effects of GH substitution (0.5–0.6 IU/kg/week) could be shown (74).

Overall, these results are very interesting, but some of them are rather conflicting. It is possible that there is a difference between CO-GHD and AO-GHD which could explain the different results found by Deijen and Baum. It can be speculated that the occurrence of GHD during a more vulnerable phase or the longer duration of GHD in CO-GHD or a combination of both of these leads to more pronounced cognitive deficits in CO-GHD than in AO-GHD patients.

Another possible explanation is that the effects seen depend on the dose of GH. It is quite striking that in the study of Baum *et al.* (73), GH doses were much lower than in all other studies cited above. Baum's study was the only one in which the GH doses were adjusted to IGF-I levels. In all the other studies the applied doses were three to ten times higher, which often led to supraphysiological IGF-I levels. In the placebo-controlled phase, Deijen *et al.* (68) could only find cognitive improvement with doses which led to supranormal IGF-I levels, and in the open-label phase a placebo effect cannot be excluded. Degerblad *et al.* (74) found no effects even though they used a very high dose. It is possible that either GH does not exert an effect on the assessed cognitive functions or that the duration and the power of the study were too low to measure any existing effects. Deijen *et al.* (68) found improvements in learning and recognition but, like Oertel *et al.* (72), no changes in short-term and long-term verbal memory. Taking the above-mentioned data together, it is quite possible that effects on cognitive functions can only be achieved with supraphysiological GH doses. On the other hand, it cannot be ruled out that in some studies the duration of the study was too short to show any effects. There is still a need for placebo-controlled trials with large case numbers in which the effects of physiological GH doses on different aspects of cognition, sex and etiology of GHD can be studied.

### **Growth hormone, insulin-like growth factor-I and cognition during aging**


In the elderly, GH secretion and serum IGF-I levels decrease (75–77) and there is a reduced expression of GH receptors in the brain (7). Normal aging is accompanied by slight decreases in some aspects of cognitive function (78–80). In patients with dementia, decreased serum IGF-I levels can be found (81, 82). It has been a subject of debate as to whether there is a

**Table 1** Influence of GH substitution on cognitive function in GH-deficient patients.

Reference	Date	n	Dose (IU/kg/week)	Design	Results
Almqvist <i>et al.</i> (69)	1986	5	~0.34	Adults with CO-GHD, DBPC, cross-over GH for 4 weeks	Improvement in face recognition
Degerblad <i>et al.</i> (74)	1990	6	0.5–0.6	DBPC, cross-over GH for 12 weeks in GHD	No objective change in cognitive function
Sartorio <i>et al.</i> (70)	1995	8	0.5	Adults with CO-GHD, open-label GH for 6 months	Improvement in symbol-number association (cognitive, visual-motoric and procedural speed)
Deijen <i>et al.</i> (68)	1998	48	~0.175, 0.35 or 0.525	Men with CO-GHD: 6 months PC with 3 different doses, 18 months open-label	In first 6 months improvement of memory (associate learning task, associate recognition task) compared with placebo only with supraphysiological doses. After 1 year improvement in all treatment groups
Baum <i>et al.</i> (73)	1998	40	0.042–0.126, adjusted to IGF-I	Men with AO-GHD: 18 months DBPC	No effect of GH on cognitive functions
Soares <i>et al.</i> (71)	1999	9	0.25 adjusted to IGF-I	DBPC 6 months GH, followed by 6 months open-label GH in adults with GHD DBPC	Improvement in attentional parameters (digit backward, verbal fluency, vocabulary, picture arrangements, comprehension)
Oertel <i>et al.</i> (72)	Unpublished data	18	0.25	GH for 6 months DBPC, followed by 6 months open-label; 16 AO-GHD, 2 CO-GHD	Improvement in attention (digit cancellation test, trail making test). No effect on verbal memory or nonverbal intelligence

The applied GH doses are shown in IU/kg/week. For Almqvist *et al.* (69) and Deijen *et al.* (68) the values were approximated. In Almqvist *et al.* (69) all probands received 8 IU GH 3 × /week independently of body weight. This corresponds to 0.34 IU/kg/week in a 70 kg proband. In Deijen *et al.* (68) 3 different doses with 1, 2, and 3 IU/m<sup>2</sup> /day were administered. The values shown were calculated for a man of 80 kg and 180 cm.

GHD, growth hormone deficiency; CO-GHD, childhood-onset GHD; AO-GHD, adulthood-onset GHD; GHRH, growth hormone releasing hormone; GH, growth hormone; IGF-I, insulin-like growth factor-I; PC, placebo-controlled; DBPC, double-blind placebo-controlled.



**Figure 2** Effect of GH replacement on measures of attention. In the digit cancellation test, the subjects were asked to search for the target letter 'd' with two dashes above and/or below it, among 'ds' with different numbers of dashes or 'ps' with two dashes. The test variable was the number of correctly cancelled items per minute. In the trail-making test subjects had to connect numbers from 1 to 90 in an ascending order. The number of items correctly connected per minute was used as the test variable. Tests were standardized to baseline. Values are shown as means  $\pm$  S.E.M., expressed in % of baseline. \* $P < 0.05$ , indicates significant improvement in the GH group compared with baseline (tests with contrasts); + $P < 0.05$ , indicates significant differences between groups (tests of simple effects). In both tests, patients showed significant improvements only under GH substitution and not under placebo treatment. After 6 months, patients under GH substitution showed significantly better scores in the digit cancellation test than patients under placebo treatment, whereas the differences between the GH and the placebo group in the trail-making test remained not significant. Data modified from (72).

relation between the decline of the somatotrophic axis and cognitive function during normal aging.

Table 2 shows an overview of the studies that have examined the relationship between cognition and the somatotrophic system during aging. In 104 elderly men, Papadakis *et al.* found a correlation between age and cognitive status, but no association of IGF-I with cognition independent of age (83). On the other hand, Morley *et al.* found a correlation between measures of cognitive function and both IGF-I and the ratio of IGF-I to GH in 65 healthy men between the ages of 20 and 84 years (84). However, in this trial, the correlation of IGF-I to cognitive function was less strong than the correlations of IGF-I to age, and cognitive function to age. It is therefore possible that both IGF-I and cognitive function only show a correlation to each other due to their close association to aging rather than being independently related. Further studies with smaller case numbers have also shown independent correlations of IGF-I and cognition (85–87). Paolisso *et al.* (87) examined the ratio of IGF-I to IGF-binding protein-3 (IGFBP-3) as IGF-I is bound mainly to IGFBP-3 in the serum and thus

IGF-I/IGFBP-3 reflects serum levels of active, bioavailable IGF-I. After adjustment for age, they found a positive correlation of both IGF-I and IGF-I/IGFBP-3 molar ratio with cognitive function in groups of elderly individuals of 75–99 years of age and in centenarians. Moreover, in a large prospective study it was found that with high levels of IGF-I as well as IGF-I/IGFBP-3 there is a low risk of cognitive decline over 2 years (89).

### **Effects of growth hormone or insulin-like growth factor-I therapy during aging**

Not many studies on the effects of GH or IGF-I treatment on cognition in normal aging subjects have been conducted. In a placebo-controlled trial, Papadakis *et al.* examined the effects of GH administration for 6 months (0.27 IU/kg/week) in healthy elderly men (mean age 75 years) with low IGF-I levels (90) and Friedlander *et al.* studied the effects of IGF-I in postmenopausal women over 60 years of age for one year in a placebo-controlled study (91). In both trials, no effect

**Table 2** GH, IGF-I and cognitive function in the elderly.

Reference	Date	n	Design	Results
Papadakis <i>et al.</i> (83)	1995	104	Cross-sectional, healthy men 74–94 years	No independent correlation between IGF-I and cognitive function
Morley <i>et al.</i> (84)	1997	65	Cross-sectional, healthy men 20–84 years	Correlation of IGF-I and IGF-I/GH with age and cognitive parameters, negative correlation of IGF-I with age
Paolisso <i>et al.</i> (88)	1997	96	Cross-sectional, healthy probands $n = 30$ 21–49 years, $n = 30$ 75–99 years, $n = 19$ centenarians	IGF-I/IGFBP-3 higher in centenarians than in probands 75–99 years of age. Independent correlation of IGF-I and IGF-I/IGFBP-3 with MMSE in centenarians and in probands 75–99 years of age
Rollero <i>et al.</i> (85)	1998	22	Cross-sectional, healthy probands 65–86 years	Independent correlation of MMSE with IGF-I
Aleman <i>et al.</i> (86)	1999	25	Cross-sectional, healthy men 65–76 years	Independent correlation of IGF-I with digit symbol test and concept shifting task
Aleman <i>et al.</i> (87)	2000	17	Cross-sectional, healthy men 66–76 years	Negative correlation of GH secretory capacity (GHRH-GHRP-6 test) and positive correlation of IGF-I with cognitive function
Kalmijn <i>et al.</i> (89)	2000	186	Prospective over 1.9 years, healthy probands, 55–80 years	Correlation of IGF-I and IGF-I/IGFBP-3 with reduced cognitive decline
Arai <i>et al.</i> (81)	2001	49	Cross-sectional, centenarians	Higher prevalence of dementia in persons with low IGF-I
Murialdo <i>et al.</i> (82)	2001	25	Cross-sectional, 25 patients with Alzheimer's disease, 12 control persons	IGF-I decreased in Alzheimer's disease
Papadakis <i>et al.</i> (90)	1996	52	DBPC, healthy men over 69 years (mean age 75 years), GH for 6 months	No effect of GH on cognitive abilities
Friedlander <i>et al.</i> (90)	2001	16	DBPC, IGF-I for 12 months in postmenopausal women over 60 years	No effect of IGF-I on mood and memory

GHRH, growth hormone releasing hormone; GHRP-6, growth hormone releasing peptide-6; GH, growth hormone; IGF-I, insulin-like growth factor-I; IGFBP-3, IGF-binding protein-3; DBPC, double-blind placebo-controlled; MMSE, mini mental state examination.

on mood, memory or other aspects of cognitive function could be found.

Overall, there is some evidence that the somatotrophic axis may have an influence on cognitive abilities even in the physiological range, although there are probably many other factors. However, a positive effect of GH or IGF-I therapy on cognitive function in the elderly has not been shown. The underlying mechanisms for the decline in IGF-I seen in GHD and normal aging are not the same and the effects of replacement therapy may, therefore, be different. It is possible that the lack of effectiveness of GH application in the elderly is due to the fact that during aging not only do GH secretion and IGF-I levels decline but so does GH receptor expression in the brain, thus reducing central sensitivity to GH. Moreover, the fact that IGF-I is positively correlated with cognitive function, and that low IGF-I levels are found in dementia does not necessarily imply that low IGF-I levels are the reason for impaired cognitive function. And finally, it should be considered that during normal and pathological aging changes in other hormonal systems such as a decline in dehydroepiandrosterone (DHEA), DHEA sulfate, and sex steroids and an increase in nocturnal cortisol levels, may occur. These hormones may also exert neurotropic actions and additionally may influence cognitive changes during aging. For the present, it cannot be ruled out that low IGF-I is simply a side-effect of cognitive decline without any functional significance. Further studies are needed.

## Quality of life

### Growth hormone deficiency

Adults who were substituted with GH for childhood GHD showed impaired emotional status, physical mobility and social isolation after discontinuation of GH replacement (92). Reduced quality of life (QoL), assessed with the quality of life-assessment in growth hormone deficiency in adults (QOL-AGHDA), a questionnaire specifically designed for evaluation of QoL in GHD, was found in a study of more than 900 untreated GH-deficient patients (93). In a prospective trial over two years it was found that patients showed a decline in general scores of QoL and higher scores of depression and anxiety after termination of GH substitution (93). On the other hand, in other studies it could be shown that the degree of impairment of QoL is also dependent on the functioning of other pituitary axes (67, 95). GH-deficient subjects showed an increased incidence of social phobia (96), depression and fatigue (97) and decreased self-esteem and QoL in self-rating questionnaires (97–99). In 86 patients with AO-GHD worse scores for energy, social isolation, emotional reaction and sex life were described in the Nottingham Health Profile (NHP) (99). On the other hand, in a study of 48 patients with pituitary disease with at least two

hormone deficiencies and low IGF-I, no differences compared with controls were found in the assessment of QoL using two questionnaires, the Short Form 36 and the General Well-Being Schedule (100). It has been found that patients with AO-GHD show higher impairments in QoL than patients with CO-GHD (101, 102).

### Effects of growth hormone replacement

Improvement in QoL after GH substitution in GH-deficient subjects has been found in an open-label study as assessed with the NHP and the Psychological General Well-Being Schedule (PGWB) (103) and in several placebo-controlled trials with the NHP (101, 104–106), the Comprehensive Psychological Rating Scale (107), the Kellner Symptom Questionnaire (108), and the Hopkins Symptom Checklist (HSCL) (104). However, other placebo-controlled trials showed no impact of GH substitution on QoL assessed with the NHP, the PGWB and the Minnesota Multiphasic Personality Inventory-2 (73), the HSCL and the Profile of Mood States (68) and the Disease Specific Questionnaire, the Symptom Checklist-90 and the Social Adjustment Scale (109). There are several possible reasons for these different findings. First, not all authors used the same questionnaires. Different methods may have led to different results. As mentioned above the GH dose given in the study of Baum *et al.* (73) was lower than in the other studies. In the study of Florkowski *et al.* (109) a cross-over design with a three-months active period was chosen and it is possible that this period was too short to produce measurable effects. Indeed, many studies showed continuous improvement in QoL over 1 or 2 years (101, 105). On the other hand, in these studies, both of which were placebo-controlled for the first 6 months and open-label afterwards, in some aspects, positive findings could only be shown after a longer period. Thus, it cannot be excluded that some of the positive findings are simply due to a placebo effect. A further reason for negative findings could be a lack of specificity of the questionnaires used.

As a consequence, specific questionnaires to evaluate successful treatment of GH substitution have been developed. The questionnaires existing at the moment are the modified impact scale (IS), the modified life fulfillment scale (LFS) (110), the growth hormone deficiency questionnaire (GHDQ) (106), the QOL-AGHDA (111), and the Questions on Life Satisfaction Modules-Hypopituitarism (QLS<sup>M</sup>-H) (112).

Interestingly, in one trial, placebo-controlled for 6 months and open-label for another 6 months, that evaluated the effectiveness of GH substitution using the NHP and the GHDQ, significant improvements were seen in some aspects of the NHP but no change in scores were seen for the GHDQ (106). The GHDQ, which was specifically developed for that trial, consisted of 30 questions related to the three subscales of energy,

mood, and sleep, each answered on a 10-cm visual analog scale on a portable computer.

The LFS is a self-assessment questionnaire consisting of 12 items divided into two subscales: personal fulfillment and material fulfillment. It consists of two questionnaires, the first one dealing with the importance of and the second one dealing with satisfaction with various aspects of life. The IS consists of ten items dealing with the impact the disease has on different aspects of life. Both questionnaires have been tested for reliability and validity (110) and used in a 12-month trial with a 6-month phase being placebo-controlled and another 6-month phase being open-label for the efficacy of GH replacement in a group of 32 patients (105). Here, no significant change in the LFS but an improvement in the IS was shown.

The QOL-AGHDA is the best validated questionnaire at the present time (102, 113–120). It is a questionnaire containing 25 items that can be answered dichotomously ('yes' or 'no'). The score is calculated by adding all the items answered with 'yes', with a high score indicating poor QoL. It exists in five languages and has shown good reliability and validity in each language version. Moreover, it is used in an international database monitoring the effects of GH replacement (111).

The QLS<sup>M</sup>-H consists of nine items. Each item can be rated according to its importance to the individual as well as the individual's satisfaction with it on a 5-point scale, with low scores implicating low QoL. The questionnaire was designed to have a high specificity for hypopituitarism and, additionally, provides the opportunity to weight each item according to its relative importance to the individual. In an open-label trial with 717 GHD patients and 2700 control individuals it showed a high reliability and validity and significant improvements in QoL during GH substitution (112).

Taken together, data on the effects of GH on QoL are still controversial. If GH substitution has an effect on QoL, the LFS and the the GHDQ appear to lack the sensitivity to measure it. The IS has shown improvements in a placebo-controlled trial, but it has not been studied in a large sample. The QOL-AGHDA has been used in many trials and in a large database and the QLS<sup>M</sup>-H has also been tested with a large population. However, both of them lack results from placebo-controlled studies. This is particularly important, as a subjective measure such as QoL is highly susceptible to a placebo effect. Therefore, testing of these questionnaires in long-term, placebo-controlled studies is needed to evaluate their real usefulness.

## Mood and depression

Healthy young men with high depression and anxiety scores show a reduced or no increase in GH secretion after physical exercise (121). In depressive patients, reduced nocturnal and 24-h GH secretion have been

found (122–124), but other authors have found unchanged (125) or even increased 24-h GH secretion due to elevated daytime levels (126, 127). The GH response to stimulation testing is reduced (122, 128) but elevated levels of IGF-I have been observed (125, 129). In depressed children, a decreased response to dynamic testing has also been found (130, 131). In addition, in children and adolescents who are not depressive but have a high genetic risk of depression, a reduced increase in GH in response to stimulation testing was found (132).

Since well-being appears to be affected by GH and the somatotrophic axis is altered in mood disorders, it can be speculated that GH or IGF-I affect brain areas or neurotransmitter systems that are related to mood and well-being. In fact, an increase in beta-endorphin in the CSF could be shown after GH substitution in GH-deficient adults (13), and in rats after intracerebroventricular GH administration (9). Beta-endorphin is an endogenous opioid that can stimulate dopamine secretion in the brain reward system (133, 134). In the mammalian brain, this consists of synaptically interconnected neurons associated with the medial forebrain bundle, linking the ventral tegmental area, nucleus accumbens, and ventral pallidum. Electrical stimulation of this circuit supports intense self-stimulation in animals and, in humans, produces intense pleasure or euphoria. This circuit is strongly implicated in the neural substrates of drug addiction as well as in the pleasures produced by natural rewards such as food and sex (135). Burman *et al.* found no change in beta-endorphin in the CSF after GH substitution but a decrease in free thyroxine (T4) and homovanillic acid (HVA) and an increase in aspartate levels (14, 57). A reduction in HVA levels in the CSF can also be seen after successful treatment of depression with antidepressant agents. HVA reflects the dopamine metabolism in the brain whereas aspartate, as discussed above, is a ligand of the NMDA receptor that has been implicated in memory (58) and attentional function (59) of the mammalian brain. A possible reason for decreased T4 is an increased conversion into triiodothyronine in the brain (57).

## Sleep

GH secretion is associated with sleep rhythm. In young men the GH peak at sleep onset is normally the most reproducible peak. In middle-aged and elderly men this is often the only phase during which measurable GH secretion takes place. In premenopausal women the GH peak at sleep onset also occurs but does not normally constitute the main part of the 24-h secretion. In shifts of sleep onset the GH peak still normally occurs at sleep onset (136), but in about 25% of young men a GH peak occurs before sleep onset (137). The somatotrophic axis interacts with sleep on many levels: there is a close relationship between the

**Table 3** Influence of the somatotrophic system on sleep and hormone secretion during sleep.

Hormone administered	Reference	SWS	REM sleep	GH	Cortisol
GHRH	146–148	↑	↑/–	↑	↓
GH	142, 143	↑/–	↑/–	↑/n.a.	n.a.
GH-secretagoges	153–156	↑ (Ghrelin, MK-677)–(GHRP-6)	↑ (MK-677) – (Ghrelin, GHRP-6)	– (MK-677) ↑ (Ghrelin, GHRP-6)	– (MK-677) ↑ (Ghrelin, GHRP-6)
Somatostatin	146, 150, 151	↓ (old pr.), – (young pr.)	↓ (old pr.), – (young pr.)	–	n.a.
CRH	157, 159	↓	↓	↑	↑
ACTH	158	–	↓	↑	↑
Cortisol	158, 160	↑	↓	↑	↑

↑, increase; ↓, decrease; –, unchanged; pr., probands; n.a., not assessed; SWS, slow wave sleep; GHRP-6, growth hormone releasing peptide-6; CRH, corticotropin releasing hormone; ACTH, adrenocorticotropin.

occurrence of slow wave sleep (SWS) of the sleep phases III and IV, and pulsatile GH secretion. In addition, with increasing age, a parallel exponential decline in SWS duration and nocturnal GH secretion occurs (138). Therefore, a common cause for SWS and the regulation of pulsatile GH secretion has been assumed (139). In Table 3, an overview of the effects of different components of the somatotrophic axis and its functional antagonists on sleep structure is shown. In patients with isolated GHD an increase in total sleep time and reduced SWS have been found (140, 141). In healthy individuals, GH has been found to reduce SWS and increase rapid eye movement (REM) sleep duration (142), or to exert no changes in sleep pattern (143). In GH-deficient adults, on the other hand, increases in SWS and REM sleep have been found after GH substitution (144), whereas in children with GHD a decrease in stage III sleep after GH replacement has been described (145).

When GHRH is applied in a pulsatile fashion during the first half of the night, an increase in SWS and GH secretion and a suppression in cortisol can be seen (146). Other authors found enhanced REM sleep after a bolus injection of GHRH but an increase in SWS only after sleep deprivation or when GHRH was given in the third REM sleep period (147). When comparing pulsatile GHRH injections to continuous GHRH infusions, a more pronounced effect of pulsatile GHRH administration on sleep promotion and GH secretion was found (148). During aging the response of GH and SWS to GHRH declines (149). Somatostatin decreases SWS in elderly subjects (150), whereas in young individuals no effect of somatostatin on sleep EEG can be found (146, 151). In the young it is possible that GHRH tonus is high enough to antagonize the sleep disturbing effects of somatostatin, while this effect vanishes with a decrease in GHRH tonus due to aging.

In normal and hypophysectomized rats GHRH stimulates non-REM sleep. REM sleep is also stimulated by GHRH in normal rats but not in hypophysectomized rats (152). Thus non-REM sleep appears to be directly stimulated by GHRH, whereas REM sleep seems to be mediated by GH but not by GHRH. The fact that GHRH stimulates non-REM sleep both in normal and hypophysectomized rats implies that GHRH directly stimulates non-REM sleep. On the other hand, both GHRH and GH increase REM sleep and GH reduces SWS which implies that REM sleep is stimulated by GH (139) and that the decrease in SWS by GH is induced by a negative feedback mechanism on GHRH. Pulsatile administration of ghrelin, the natural GH secretagog receptor ligand, leads to enhanced SWS and stimulation of GH and cortisol secretion (153). A single dose of GH releasing peptide (RP)-6, a synthetic GH secretagog, induces an increase in phase II sleep (154, 155). Oral administration of MK-677, another GH secretagog, over one week leads to an increase in REM and phase IV sleep (156).

In some respects the adrenocorticotrophic system appears to act as an antagonist of the somatotrophic system. Corticotropin releasing hormone (CRH) reduces nocturnal GH secretion (157), and CRH, adrenocorticotropin and cortisol decrease REM sleep duration (158); SWS duration is reduced by CRH (159) and increased by cortisol (158, 160). This implies a central inhibition of SWS by CRH and an inhibition of REM sleep by cortisol. During normal aging and in depression sleep becomes shallow, SWS and the activity of the somatotrophic system decreases, and the activity of the adrenocorticotrophic system increases, so that the balance between the two systems shifts towards the adrenocorticotrophic system (139).

In summary, these data show that neuropeptides, particularly GHRH and CRH play an important role in the regulation of sleep and act as functional antagonists. However, it has also been shown that both GH and cortisol can also exert direct effects on sleep structure.

## Conclusions

Data on the effects of GH, IGF-I and GHRH on brain function have been accumulating in recent years, but understanding is still lacking as to exactly how GHRH, GH, and IGF-I interact with the brain. Putative mechanisms include local production of hormones, transport via the BBB, influence on levels of neurotransmitters, and the regulation of genes such as BDNF and c-fos.

Both GH and IGF-I appear to exert neuroprotective effects in pathological brain damage, as well as in normal brain aging. However, most results have been obtained in animal studies and evidence for a possible clinical application is scarce. Cognitive impairments can be seen in GHD and some studies have shown improvement after GH replacement. As some studies have produced conflicting results, further evaluation is needed. An impact of GH on mood and well-being has been found with GH-deficient patients showing a reduced quality of life and a reduced sense of well-being. Specific questionnaires that have been developed to assess the efficacy of GH replacement still need to be evaluated in placebo-controlled trials. In depression, a changed GH secretion has been found, the meaning of which still remains to be elucidated. Both GHRH and GH seem to influence sleep structure and sleep is altered in GHD.

## Acknowledgements

We are grateful to Prof. Dr A Steiger and Dr U Renner for critically reading the manuscript and for helpful comments.

## References

- Salomon F, Cuneo RC, Hesp R & Sonksen PH. The effects of treatment with recombinant human growth hormone on body composition and metabolism in adults with growth hormone deficiency. *New England Journal of Medicine* 1989 **321** 1797–1803.
- Jorgensen JO, Pedersen SA, Thuesen L, Jorgensen J, Ingemann-Hansen T, Skakkebaek NE *et al.* Beneficial effects of growth hormone treatment in GH-deficient adults. *Lancet* 1989 **1** 1221–1225.
- Kann P, Piepkorn B, Schehler B, Andreas J, Lotz J, Prellwitz W *et al.* Effect of long-term treatment with GH on bone metabolism, bone mineral density and bone elasticity in GH-deficient adults. *Clinical Endocrinology* 1998 **48** 561–568.
- Kann P, Piepkorn B, Schehler B, Lotz J, Prellwitz W & Beyer J. Growth hormone substitution in growth hormone-deficient adults: effects on collagen type I synthesis and skin thickness. *Experimental and Clinical Endocrinology and Diabetes* 1996 **104** 327–333.
- Gotherstrom G, Svensson J, Koranyi J, Alpsten M, Bosaeus I, Bengtsson BA *et al.* A prospective study of 5 years of GH replacement therapy in GH-deficient adults: sustained effects on body composition, bone mass, and metabolic indices. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 4657–4665.
- Wüster C, Abs R, Bengtsson BA, Benmarker H, Feldt-Rasmussen U, Hernberg-Stahl E *et al.* The influence of growth hormone deficiency, growth hormone replacement therapy, and other aspects of hypopituitarism on fracture rate and bone mineral density. *Journal of Bone and Mineral Research* 2001 **16** 398–405.
- Lai Z, Roos P, Zhai O, Olsson Y, Fholenhag K, Larsson C *et al.* Age-related reduction of human growth hormone-binding sites in the human brain. *Brain Research* 1993 **621** 260–266.
- Nyberg F & Burman P. Growth hormone and its receptors in the central nervous system – location and functional significance. *Hormone Research* 1996 **45** 18–22.
- Nyberg F. Growth hormone in the brain: characteristics of specific brain targets for the hormone and their functional significance. *Frontiers in Neuroendocrinology* 2000 **21** 330–348.
- Castro JR, Costoya JA, Gallego R, Prieto A, Arce VM & Senaris R. Expression of growth hormone receptor in the human brain. *Neuroscience Letters* 2000 **281** 147–150.
- Rosenfeld RG & Bengtsson BA. Effects of growth hormone and insulin-like growth factors on the central nervous system. *Acta Paediatrica* 1994 **406** (Suppl) 89–91.
- Rolandi E, Perria C, Cicchetti V, Sannia A, Magnani G, Rivano C *et al.* Pituitary hormone concentrations in cerebrospinal fluid in patients with prolactin and growth hormone-secreting tumors. *Journal of Neurosurgical Sciences* 1982 **26** 173–178.
- Johansson JO, Larson G, Andersson M, Elmgren A, Hynsjo L, Lindahl A *et al.* Treatment of growth hormone-deficient adults with recombinant human growth hormone increases the concentration of growth hormone in the cerebrospinal fluid and affects neurotransmitters. *Neuroendocrinology* 1995 **61** 57–66.
- Burman P, Hetta J, Wide L, Mansson JE, Ekman R & Karlsson FA. Growth hormone treatment affects brain neurotransmitters and thyroxine. *Clinical Endocrinology* 1996 **44** 319–324.
- Hojvat S, Baker G, Kirsteins L & Lawrence AM. Growth hormone (GH) immunoreactivity in the rodent and primate CNS: distribution, characterization and presence posthypophysectomy. *Brain Research* 1982 **239** 543–557.
- Mustafa A, Nyberg F, Bogdanovic N, Islam A, Roos P & Adem A. Somatogenic and lactogenic binding sites in rat brain and liver: quantitative autoradiographic localization. *Neuroscience Research* 1994 **20** 257–263.
- Coculescu M. Blood-brain barrier for human growth hormone and insulin-like growth factor-I. *Journal of Pediatric Endocrinology and Metabolism* 1999 **12** 113–124.

- 18 Walsh RJ, Slaby FJ & Posner BI. A receptor-mediated mechanism for the transport of prolactin from blood to cerebrospinal fluid. *Endocrinology* 1987 **120** 1846–1850.
- 19 Lobie PE, Zhu T, Graichen R & Goh EL. Growth hormone, insulin-like growth factor I and the CNS: localization, function and mechanism of action. *Growth Hormone and IGF Research* 2000 **10** (Suppl B) S51–S56.
- 20 Adem A, Jossan SS, d'Argy R, Gillberg PG, Nordberg A, Winblad B *et al.* Insulin-like growth factor 1 (IGF-1) receptors in the human brain: quantitative autoradiographic localization. *Brain Research* 1989 **503** 299–303.
- 21 Carro E, Nunez A, Busiguina S & Torres-Aleman I. Circulating insulin-like growth factor I mediates effects of exercise on the brain. *Journal of Neuroscience* 2000 **20** 2926–2933.
- 22 Anlar B, Sullivan KA & Feldman EL. Insulin-like growth factor-I and central nervous system development. *Hormone and Metabolic Research* 1999 **31** 120–125.
- 23 Matsubara S, Sato M, Mizobuchi M, Niimi M & Takahara J. Differential gene expression of growth hormone (GH)-releasing hormone (GRH) and GRH receptor in various rat tissues. *Endocrinology* 1995 **136** 4147–4150.
- 24 Van Cauter E, Plat L & Copinschi G. Interrelations between sleep and the somatotrophic axis. *Sleep* 1998 **21** 553–566.
- 25 Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H & Kangawa K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature* 1999 **402** 656–660.
- 26 Mayo KE, Godfrey PA, Suhr ST, Kulik DJ & Rahal JO. Growth hormone-releasing hormone: synthesis and signaling. *Recent Progress in Hormone Research* 1995 **50** 35–73.
- 27 Brazeau P, Vale W, Burgus R, Ling N, Butcher M, Rivier J *et al.* Hypothalamic polypeptide that inhibits the secretion of immunoreactive pituitary growth hormone. *Science* 1973 **179** 77–79.
- 28 Vale W, Brazeau P, Rivier C, Brown M, Boss B, Rivier J *et al.* Somatostatin. *Recent Progress in Hormone Research* 1975 **31** 365–397.
- 29 Kamegai J, Tamura H, Shimizu T, Ishii S, Sugihara H & Wakabayashi I. Central effect of ghrelin, an endogenous growth hormone secretagogue, on hypothalamic peptide gene expression. *Endocrinology* 2000 **141** 4797–4800.
- 30 Kamegai J, Tamura H, Shimizu T, Ishii S, Sugihara H & Oikawa S. Regulation of the ghrelin gene: growth hormone-releasing hormone upregulates ghrelin mRNA in the pituitary. *Endocrinology* 2001 **142** 4154–4157.
- 31 Date Y, Murakami M, Kojima M, Kuroiwa T, Matsukura S, Kangawa K *et al.* Central effects of a novel acylated peptide, ghrelin, on growth hormone release in rats. *Biochemical and Biophysical Research Communications* 2000 **275** 477–480.
- 32 Peino R, Baldelli R, Rodriguez-Garcia J, Rodriguez-Segade S, Kojima M, Kangawa K *et al.* Ghrelin-induced growth hormone secretion in humans. *European Journal of Endocrinology* 2000 **143** R11–R14.
- 33 Takaya K, Ariyasu H, Kanamoto N, Iwakura H, Yoshimoto A, Harada M *et al.* Ghrelin strongly stimulates growth hormone release in humans. *Journal of Clinical Endocrinology and Metabolism* 2000 **85** 4908–4911.
- 34 Arvat E, Maccario M, Di Vito L, Broglio F, Benso A, Gottero C *et al.* Endocrine activities of ghrelin, a natural growth hormone secretagogue (GHS), in humans: comparison and interactions with hexarelin, a nonnatural peptidyl GHS, and GH-releasing hormone. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 1169–1174.
- 35 Hataya Y, Akamizu T, Takaya K, Kanamoto N, Ariyasu H, Saijo M *et al.* A low dose of ghrelin stimulates growth hormone (GH) release synergistically with GH-releasing hormone in humans. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 4552–4555.
- 36 Wren AM, Small CJ, Abbott CR, Dhillo WS, Seal I, Cohen MA *et al.* Ghrelin causes hyperphagia and obesity in rats. *Diabetes* 2000 **50** 2540–2547.
- 37 Wren AM, Small CJ, Ward HL, Murphy KG, Dakin CL, Taheri S *et al.* The novel hypothalamic peptide ghrelin stimulates food intake and growth hormone secretion. *Endocrinology* 2000 **141** 4325–4328.
- 38 Tschöp M, Smiley DL & Heiman ML. Ghrelin induces adiposity in rodents. *Nature* 2000 **407** 908–913.
- 39 Wren AM, Seal LJ, Cohen MA, Brynes AE, Frost GS & Murphy KG. Ghrelin enhances appetite and increases food intake in humans. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 5992–5995.
- 40 Tschöp M, Weyer C, Tataranni PA, Devanarayan V, Ravussin E & Heiman ML. Circulating ghrelin levels are decreased in human obesity. *Diabetes* 2001 **50** 707–709.
- 41 Ariyasu H, Takaya K, Tagami T, Ogawa Y, Hosoda K, Akamizu T *et al.* Stomach is a major source of circulating ghrelin, and feeding state determines plasma ghrelin-like immunoreactivity levels in humans. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 4753–4758.
- 42 Caregaro L, Favaro A, Santonastaso P, Alberino F, Di Pascoli L, Nardi M *et al.* Insulin-like growth factor 1 (IGF 1), a nutritional marker in patients with eating disorders. *Clinical Nutrition* 2001 **20** 251–257.
- 43 Maccario M, Grottoli S, Aimaretti G, Gianotti L, Oleandri E, Procopio M *et al.* IGF-I levels in different conditions of low somatotrope secretion in adulthood: obesity in comparison with GH deficiency. *Panminerva Medica* 1998 **40** 98–102.
- 44 Maccario M, Ramunni J, Oleandri SE, Procopio M, Grottoli S, Savio P *et al.* Relationships between IGF-I and age, gender, body-mass, fat distribution, metabolic and hormonal variables in obese patients. *International Journal of Obesity Related Metabolic Disorders* 1999 **23** 612–618.
- 45 Chen L, Lund PK, Burgess SB, Rudisch BE & McIlwain DL. Growth hormone, insulin-like growth factor I, and motoneuron size. *Journal of Neurobiology* 1997 **32** 202–212.
- 46 Tsitouras PD, Zhong YG, Spungen AM & Bauman WA. Serum testosterone and growth hormone/insulin-like growth factor-I in adults with spinal cord injury. *Hormone and Metabolic Research* 1995 **27** 287–292.
- 47 Scheepens A, Sirimanne ES, Breier BH, Clark RG, Gluckman PD & Williams CE. Growth hormone as a neuronal rescue factor during recovery from CNS injury. *Neuroscience* 2001 **104** 677–687.
- 48 Niblock MM, Brunso-Bechtold JK & Riddle DR. Insulin-like growth factor I stimulates dendritic growth in primary somatosensory cortex. *Journal of Neuroscience* 2000 **20** 4165–4176.
- 49 Beck KD, Powell-Braxton L, Widmer HR, Valverde J & Hefti F. IGF1 gene disruption results in reduced brain size, CNS hypomyelination, and loss of hippocampal granule and striatal parvalbumin-containing neurons. *Neuron* 1995 **14** 717–730.
- 50 Dore S, Kar S, Zheng WH & Quirion R. Rediscovering good old friend IGF-I in the new millennium: possible usefulness in Alzheimer's disease and stroke. *Pharmaceutica Acta Helvetica* 2000 **74** 273–280.
- 51 Lichtenwalner RJ, Forbes ME, Bennett SA, Lynch CD, Sonntag WE & Riddle DR. Intracerebroventricular infusion of insulin-like growth factor-I ameliorates the age-related decline in hippocampal neurogenesis. *Neuroscience* 2001 **107** 603–613.
- 52 Liu XF, Fawcett JR, Thorne RG, DeFor TA & Frey WH. Intranasal administration of insulin-like growth factor-I bypasses the blood-brain barrier and protects against focal cerebral ischemic damage. *Journal of the Neurological Sciences* 2001 **187** 91–97.
- 53 Liu XF, Fawcett JR, Thorne RG & Frey WH. Non-invasive intranasal insulin-like growth factor-I reduces infarct volume and improves neurologic function in rats following middle cerebral artery occlusion. *Neuroscience Letters* 2001 **308** 91–94.
- 54 Trejo JL, Carro E & Torres-Aleman I. Circulating insulin-like growth factor I mediates exercise-induced increases in the

- number of new neurons in the adult hippocampus. *Journal of Neuroscience* 2001 **21** 1628–1634.
- 55 Carro E, Trejo JL, Busiguina S & Torres-Aleman I. Circulating insulin-like growth factor I mediates the protective effects of physical exercise against brain insults of different etiology and anatomy. *Journal of Neuroscience* 2001 **21** 5678–5684.
- 56 Alvarez XA & Cacabelos R. Effects of GRF (1–29) NH<sub>2</sub> on short-term memory: neuroendocrine and neuropsychological assessment in healthy young subjects. *Methods and Findings in Experimental and Clinical Pharmacology* 1990 **12** 493–499.
- 57 Burman P, Hetta J & Karlsson A. Effect of growth hormone on brain neurotransmitters. *Lancet* 1993 **342** 1492–1493.
- 58 Newcomer JW & Krystal JH. NMDA receptor regulation of memory and behavior in humans. *Hippocampus* 2001 **11** 529–542.
- 59 Turchi J & Sarter M. Bidirectional modulation of basal forebrain N-methyl-D-aspartate receptor function differentially affects visual attention but not visual discrimination performance. *Neuroscience* 2001 **104** 407–417.
- 60 Sartorio A, Conti A, Molinari E, Riva G, Morabito F & Faglia G. Growth, growth hormone and cognitive functions. *Hormone Research* 1996 **45** 23–29.
- 61 Abbott D, Rotnem D, Genel M & Cohen DJ. Cognitive and emotional functioning in hypopituitary short-statured children. *Schizophrenia Bulletin* 1982 **8** 310–319.
- 62 Steinhausen HC & Stahnke N. Negative impact of growth-hormone deficiency on psychological functioning in dwarfed children and adolescents. *European Journal of Pediatrics* 1977 **126** 263–270.
- 63 Rikken B, van Busschbach J, le Cessie S, Manten W, Spermon T, Grobbee R *et al.* Impaired social status of growth hormone deficient adults as compared to controls with short or normal stature. Dutch Growth Hormone Working Group. *Clinical Endocrinology* 1996 **44** 489–490.
- 64 Dean HJ, McTaggart TL, Fish DG & Friesen HG. The educational, vocational, and marital status of growth hormone deficient adults treated with growth hormone during childhood. *American Journal of Diseases of Children* 1985 **139** 1105–1110.
- 65 Peace KA, Orme SM, Thompson AR, Padayatti S, Ellis AW & Belch PE. Cognitive dysfunction in patients treated for pituitary tumors. *Journal of Clinical and Experimental Neuropsychology* 1997 **19** 1–6.
- 66 Bülow B, Hagmar L, Orbaek P, Österberg K & Erfurth EM. High incidence of mental disorders, reduced mental well-being and cognitive dysfunction in hypopituitary women with GH deficiency treated for pituitary disease. *Clinical Endocrinology* 2002 **56** 183–193.
- 67 Deijen JB, de Boer H, Blok GJ & van der Veen EA. Cognitive impairments and mood disturbances in growth hormone deficient men. *Psychoneuroendocrinology* 1996 **21** 313–322.
- 68 Deijen JB, de Boer H & van der Veen EA. Cognitive changes during growth hormone replacement in adult men. *Psychoneuroendocrinology* 1998 **23** 45–55.
- 69 Almqvist O, Thoren M, Saaf M & Eriksson O. Effects of growth hormone substitution on mental performance in adults with growth hormone deficiency: a pilot study. *Psychoneuroendocrinology* 1986 **11** 347–352.
- 70 Sartorio A, Molinari E, Riva G, Conti A, Morabito F & Faglia G. Growth hormone treatment in adults with childhood onset growth hormone deficiency: effects on psychological capabilities. *Hormone Research* 1995 **44** 6–11.
- 71 Soares CN, Musolino NR, Cunha NM, Caires MA, Rosenthal MC, Camargo CP *et al.* Impact of recombinant human growth hormone (RH-GH) treatment on psychiatric, neuropsychological and clinical profiles of GH deficient adults. A placebo-controlled trial. *Arquivos de Neuro-Psiquiatria* 1999 **57** 182–189.
- 72 Oertal H, Schneider HJ, Stalla GK, Holsboer F & Zihl J. The effect of growth hormone substitution of cognitive performance in adult patients with hypopituitarism. *Psychoneuroendocrinology* (In Press).
- 73 Baum HB, Katznelson L, Sherman JC, Biller BM, Hayden DL, Schoenfeld DA *et al.* Effects of physiological growth hormone (GH) therapy on cognition and quality of life in patients with adult-onset GH deficiency. *Journal of Clinical Endocrinology and Metabolism* 1998 **83** 3184–3189.
- 74 Degerblad M, Almqvist O, Grunditz R, Hall K, Kaijser L, Knutsson E *et al.* Physical and psychological capabilities during substitution therapy with recombinant growth hormone in adults with growth hormone deficiency. *Acta Endocrinologica* 1990 **123** 185–193.
- 75 Gelato MC. Aging and immune function: a possible role for growth hormone. *Hormone Research* 1996 **45** 46–49.
- 76 Rudman D, Kutner MH, Rogers CM, Lubin ME, Fleming GA & Bain RP. Impaired growth hormone secretion in the adult population: relation to age and adiposity. *Journal of Clinical Investigation* 1981 **67** 1361–1369.
- 77 Rudman D. Growth hormone, body composition, and aging. *Journal of the American Geriatric Society* 1985 **33** 800–807.
- 78 Ritchie K, Touchon J, Ledesert B, Leibovici D & Gorce AM. Establishing the limits and characteristics of normal age-related cognitive decline. *Revue d'Epidemiologie et de Santé Publique* 1997 **45** 373–381.
- 79 Compton DM, Bachman LD, Brand D & Avet TL. Age-associated changes in cognitive function in highly educated adults: emerging myths and realities. *International Journal of Geriatric Psychiatry* 2000 **15** 75–85.
- 80 Capurso A, Panza F, Solfrizzi V, Torres F, Capurso C, Mastroianni F *et al.* Age-related cognitive decline: evaluation and prevention strategy. *Recenti Progressi in Medicina* 2000 **91** 127–134.
- 81 Arai Y, Hirose N, Yamamura K, Shimizu K, Takayama M, Ebihara *et al.* Serum insulin-like growth factor-1 in centenarians: implications of IGF-1 as a rapid turnover protein. *Journals of Gerontology Series A: Biological Sciences and Medical Sciences* 2001 **56** M79–M82.
- 82 Murialdo G, Barreca A, Nobili E, Rollero A, Timossi G, Gianelli MV *et al.* Relationships between cortisol, dehydroepiandrosterone sulphate and insulin-like growth factor-I system in dementia. *Journal of Endocrinological Investigation* 2001 **24** 139–146.
- 83 Papadakis MA, Grady D, Tierney MJ, Black D, Wells L & Grunfeld C. Insulin-like growth factor 1 and functional status in healthy older men. *Journal of the American Geriatric Society* 1995 **43** 1350–1355.
- 84 Morley JE, Kaiser F, Raum WJ, Perry HM, Flood JF, Jensen J *et al.* Potentially predictive and manipulable blood serum correlates of aging in the healthy human male: progressive decreases in bioavailable testosterone, dehydroepiandrosterone sulfate, and the ratio of insulin-like growth factor 1 to growth hormone. *PNAS* 1997 **94** 7537–7542.
- 85 Rollero A, Murialdo G, Fonzi S, Garrone S, Gianelli MV, Gazzero E *et al.* Relationship between cognitive function, growth hormone and insulin-like growth factor I plasma levels in aged subjects. *Neuropsychobiology* 1998 **38** 73–79.
- 86 Aleman A, Verhaar HJ, de Haan EH, de Vries WR, Samson MM, Drent ML *et al.* Insulin-like growth factor-I and cognitive function in healthy older men. *Journal of Clinical Endocrinology and Metabolism* 1999 **84** 471–475.
- 87 Aleman A, de Vries WR, de Haan EH, Verhaar HJ, Samson MM & Koppeschaar HP. Age-sensitive cognitive function, growth hormone and insulin-like growth factor 1 plasma levels in healthy older men. *Neuropsychobiology* 2000 **41** 73–78.
- 88 Paolisso G, Ammendola S, Del Buono A, Gambardella A, Riondino M, Tagliamonte MR *et al.* Serum levels of insulin-like growth factor-I (IGF-I) and IGF-binding protein-3 in healthy centenarians: relationship with plasma leptin and lipid concentrations, insulin action, and cognitive function. *Journal of Clinical Endocrinology and Metabolism* 1997 **82** 2204–2209.

- 89 Kalmijn S, Janssen JA, Pols HA, Lamberts SW & Breteler MM. A prospective study on circulating insulin-like growth factor I (IGF-I), IGF-binding proteins, and cognitive function in the elderly. *Journal of Clinical Endocrinology and Metabolism* 2000 **85** 4551–4555.
- 90 Papadakis MA, Grady D, Black D, Tierney MJ, Gooding GA, Schambelan M *et al.* Growth hormone replacement in healthy older men improves body composition but not functional ability. *Annals of Internal Medicine* 1996 **124** 708–716.
- 91 Friedlander AL, Butterfield GE, Moynihan S, Grillo J, Pollack M, Holloway L *et al.* One year of insulin-like growth factor I treatment does not affect bone density, body composition, or psychological measures in postmenopausal women. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 1496–1503.
- 92 Bjork S, Jonsson B, Westphal O & Levin JE. Quality of life of adults with growth hormone deficiency: a controlled study. *Acta Paediatrica Scandinavica* 1989 **356** (Suppl) 55–59.
- 93 Sanmanti A, Lucas A, Hawkins F, Webb SM & Ulied A. Observational study in adult hypopituitary patients with untreated growth hormone deficiency (ODA study). Socio-economic impact and health status. Collaborative ODA (Observational GH Deficiency in Adults) Group. *European Journal of Endocrinology* 2000 **141** 481–489.
- 94 Wiren L, Johannsson G & Bengtsson BA. A prospective investigation of quality of life and psychological well-being after the discontinuation of GH treatment in adolescent patients who had GH deficiency during childhood. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 3494–3498.
- 95 Sandberg DE, MacGillivray MH, Clopper RR, Fung C, LeRoux L & Alliger DE. Quality of life among formerly treated childhood-onset growth hormone-deficient adults: a comparison with unaffected siblings. *Journal of Clinical Endocrinology and Metabolism* 1998 **83** 1134–1142.
- 96 Stabler B, Tancer ME, Ranc J & Underwood LE. Evidence for social phobia and other psychiatric disorders in adults who were growth hormone deficient during childhood. *Anxiety* 1996 **2** 86–89.
- 97 Wallymahmed ME, Foy P & MacFarlane IA. The quality of life of adults with growth hormone deficiency: comparison with diabetic patients and control subjects. *Clinical Endocrinology* 1999 **51** 333–338.
- 98 McGauley GA. Quality of life assessment before and after growth hormone treatment in adults with growth hormone deficiency. *Acta Paediatrica Scandinavica* 1989 **356** (Suppl) 70–74.
- 99 Rosen T, Wiren L, Wilhelmsen L, Wiklund I & Bengtsson BA. Decreased psychological well-being in adult patients with growth hormone deficiency. *Clinical Endocrinology* 1994 **40** 111–116.
- 100 Page RC, Hammersley MS, Burke CW & Wass JA. An account of the quality of life of patients after treatment of non-functioning pituitary tumours. *Clinical Endocrinology* 1997 **46** 401–406.
- 101 Attanasio AF, Lamberts SW, Matranga AM, Birkett MA, Bates PC, Valk NK *et al.* Adult growth hormone (GH)-deficient patients demonstrate heterogeneity between childhood onset and adult onset before and during human GH treatment. Adult Growth Hormone Deficiency Study Group. *Journal of Clinical Endocrinology and Metabolism* 1997 **82** 82–88.
- 102 Murray RD, Skillicorn CJ, Howell SJ, Lissett CA, Rahim A, Smethurst LE *et al.* Influences on quality of life in GH deficient adults and their effect on response to treatment. *Clinical Endocrinology* 1999 **51** 565–573.
- 103 Carroll PV, Littlewood R, Weissberger AJ, Bogalho P, McGauley G, Sonksen PH *et al.* The effects of two doses of replacement growth hormone on the biochemical, body composition and psychological profiles of growth hormone-deficient adults. *European Journal of Endocrinology* 1997 **137** 146–153.
- 104 Burman P, Broman JE, Hetta J, Wiklund I, Erfurth EM, Hagg E *et al.* Quality of life in adults with growth hormone (GH) deficiency: response to treatment with recombinant human GH in a placebo-controlled 21-month trial. *Journal of Clinical Endocrinology and Metabolism* 1995 **80** 3585–3590.
- 105 Wallymahmed ME, Foy P, Shaw D, Hutcheon R, Edwards RH & MacFarlane IA. Quality of life, body composition and muscle strength in adult growth hormone deficiency: the influence of growth hormone replacement therapy for up to 3 years. *Clinical Endocrinology* 1997 **47** 439–446.
- 106 Cuneo RC, Judd S, Wallace JD, Perry-Keene D, Burger H, Lim-Tio S *et al.* The Australian multicenter trial of growth hormone (GH) treatment in GH-deficient adults. *Journal of Clinical Endocrinology and Metabolism* 1998 **83** 107–116.
- 107 Bengtsson BA, Eden S, Lonn L, Kvist H, Stokland A, Lindstedt G *et al.* Treatment of adults with growth hormone (GH) deficiency with recombinant human GH. *Journal of Clinical Endocrinology and Metabolism* 1993 **76** 309–317.
- 108 Giusti M, Meineri I, Malagamba D, Cuttica CM, Fattacciu G, Menichini U *et al.* Impact of recombinant human growth hormone treatment on psychological profiles in hypopituitary patients with adult-onset growth hormone deficiency. *European Journal of Clinical Investigation* 1998 **28** 13–19.
- 109 Florkowski CM, Stevens I, Joyce P, Espiner EA & Donald RA. Growth hormone replacement does not improve psychological well-being in adult hypopituitarism: a randomized crossover trial. *Psychoneuroendocrinology* 1998 **23** 57–63.
- 110 Wallymahmed ME, Baker GA, Humphris G, Dewey M & MacFarlane IA. The development, reliability and validity of a disease specific quality of life model for adults with growth hormone deficiency. *Clinical Endocrinology* 1996 **44** 403–411.
- 111 McKenna SP, Doward LC, Alonso J, Kohlmann T, Niero M, Prieto L *et al.* The QoL-AGHDA: an instrument for the assessment of quality of life in adults with growth hormone deficiency. *Quality of Life Research* 1999 **8** 373–383.
- 112 Herschbach P, Henrich G, Strasburger CJ, Feldmeier H, Marin F, Attanasio AM & Blum WF. Development and psychometric properties of a disease-specific quality of life questionnaire for adult patients with growth hormone deficiency. *European Journal of Endocrinology* 2001 **145** 255–265.
- 113 Bengtsson BA, Abs R, Bennmarker H, Monson JP, Feldt-Rasmussen U, Hernberg-Stahl E *et al.* The effects of treatment and the individual responsiveness to growth hormone (GH) replacement therapy in 665 GH-deficient adults. KIMS Study Group and the KIMS International Board. *Journal of Clinical Endocrinology and Metabolism* 1999 **84** 3929–3935.
- 114 Abs R, Bengtsson BA, Hernberg-Stahl E, Monson JP, Tauber JP, Wilton P *et al.* GH replacement in 1034 growth hormone deficient hypopituitary adults: demographic and clinical characteristics, dosing and safety. *Clinical Endocrinology* 1999 **50** 703–713.
- 115 Bülow B & Erfurth EM. A low individualized GH dose in young patients with childhood onset GH deficiency normalized serum IGF-I without significant deterioration in glucose tolerance. *Clinical Endocrinology* 1999 **50** 45–55.
- 116 Monson JP, Abs R, Bengtsson BA, Bennmarker H, Feldt-Rasmussen U, Hernberg-Stahl E *et al.* Growth hormone deficiency and replacement in elderly hypopituitary adults. KIMS Study Group and the KIMS International Board. Pharmacology and Upjohn International Metabolic Database. *Clinical Endocrinology* 2000 **53** 281–289.
- 117 Wiren L, Whalley D, McKenna S & Wilhelmsen L. Application of a disease-specific, quality-of-life measure (QoL-AGHDA) in growth hormone-deficient adults and a random population sample in Sweden: validation of the measure by rasch analysis. *Clinical Endocrinology* 2000 **52** 143–152.
- 118 Lagrou K, Xhrouet-Heinrichs D, Massa G, Vandeweghe M, Bourguignon JP, de Schepper J *et al.* Quality of life and retrospective perception of the effect of growth hormone treatment in adult patients with childhood growth hormone deficiency. *Journal of Pediatric Endocrinology and Metabolism* 2001 **14** (Suppl) 1249–1260.

- 119 Barkan AL. The 'quality of life-assessment of growth hormone deficiency in adults' questionnaire: can it be used to assess quality of life in hypopituitarism? *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 1905–1907.
- 120 Ahmad AM, Hopkins MT, Thomas J, Ibrahim H, Fraser WD & Vora JP. Body composition and quality of life in adults with growth hormone deficiency; effects of low-dose growth hormone replacement. *Clinical Endocrinology* 2001 **54** 709–717.
- 121 Harro J, Rimm H, Harro M, Grauberg M, Karelson K & Viru AM. Association of depressiveness with blunted growth hormone response to maximal physical exercise in young healthy men. *Psychoneuroendocrinology* 1999 **24** 505–517.
- 122 Voderholzer U, Laakmann G, Wittmann R, Daffner-Bujia C, Hinz A, Haag C *et al.* Profiles of spontaneous 24-hour and stimulated growth hormone secretion in male patients with endogenous depression. *Psychiatry Research* 1993 **47** 215–227.
- 123 Fiasche R, Fideleff HL, Moiseowicz J, Frieder P, Pagano SM & Holland M. Growth hormone neurosecretory dysfunction in major depressive illness. *Psychoneuroendocrinology* 1999 **20** 727–733.
- 124 Sakkas PN, Soldatos CR, Bergiannaki JD, Paparrigopoulos TJ & Stefanis CN. Growth hormone secretion during sleep in male depressed patients. *Progress in Neuropsychopharmacology and Biological Psychiatry* 1998 **22** 467–483.
- 125 Deuschle M, Blum WF, Strasburger CJ, Schweiger U, Weber B, Korner A *et al.* Insulin-like growth factor-I (IGF-I) plasma concentrations are increased in depressed patients. *Psychoneuroendocrinology* 1997 **22** 493–503.
- 126 Mendlewicz J, Linkowski P, Kerkhofs M, Desmedt D, Golstein J, Copinschi G *et al.* Diurnal hypersecretion of growth hormone in depression. *Journal of Clinical Endocrinology and Metabolism* 1985 **60** 505–512.
- 127 Linkowski P, Mendlewicz J, Kerkhofs M, Leclercq R, Golstein J, Brasseur M *et al.* 24-hour profiles of adrenocorticotropin, cortisol, and growth hormone in major depressive illness: effect of antidepressant treatment. *Journal of Clinical Endocrinology and Metabolism* 1987 **65** 141–152.
- 128 Mokrani M, Duval F, Diep TS, Bailey PE & Macher JP. Multi-hormonal responses to clonidine in patients with affective and psychotic symptoms. *Psychoneuroendocrinology* 2000 **25** 741–752.
- 129 Franz B, Buysse DJ, Cherry CR, Gray NS, Grochocinski VJ, Frank E *et al.* Insulin-like growth factor 1 and growth hormone binding protein in depression: a preliminary communication. *Journal of Psychiatric Research* 1999 **33** 121–127.
- 130 Ryan ND, Dahl RE, Birmaher B, Williamson DE, Iyengar S, Nelson B *et al.* Stimulatory tests of growth hormone secretion in prepubertal major depression: depressed versus normal children. *Journal of the American Academy of Child and Adolescent Psychiatry* 1994 **33** 824–833.
- 131 Dahl RE, Birmaher B, Williamson DE, Dorn L, Perel J, Kaufman J *et al.* Low growth hormone response to growth hormone-releasing hormone in child depression. *Biological Psychiatry* 2000 **48** 981–988.
- 132 Birmaher B, Dahl RE, Williamson DE, Perel JM, Brent DA, Axelson DA *et al.* Growth hormone secretion in children and adolescents at high risk for major depressive disorder. *Archives of General Psychiatry* 2000 **57** 867–872.
- 133 Johansson P, Ray A, Zhou Q, Huang W, Karlsson K & Nyberg F. Anabolic androgenic steroids increase beta-endorphin levels in the ventral tegmental area in the male rat brain. *Neuroscience Research* 1997 **27** 185–189.
- 134 Spanagel R & Weiss F. The dopamine hypothesis of reward: past and current status. *Trends in Neuroscience* 1999 **22** 521–527.
- 135 Gardner EL & Vorel SR. Cannabinoid transmission and reward-related events. *Neurobiology of Disease* 1998 **5** 502–533.
- 136 Van Cauter E & Copinschi G. Interrelationships between growth hormone and sleep. *Growth Hormone and IGF Research* 2000 **10** (Suppl B) S57–S62.
- 137 Mullington J, Hermann D, Holsboer F & Pollmächer T. Age-dependent suppression of nocturnal growth hormone levels during sleep deprivation. *Neuroendocrinology* 1996 **64** 233–241.
- 138 Van Cauter E, Plat L & Copinschi G. Interrelations between sleep and the somatotrophic axis. *Sleep* 1998 **21** 553–566.
- 139 Steiger A & Holsboer F. Neuropeptides and human sleep. *Sleep* 1997 **20** 1038–1052.
- 140 Astrom C & Jochumsen PL. Decrease in delta sleep in growth hormone deficiency assessed by a new power spectrum analysis. *Sleep* 1989 **12** 508–515.
- 141 Astrom C & Lindholm J. Growth hormone-deficient young adults have decreased sleep. *Neuroendocrinology* 1990 **51** 82–94.
- 142 Mendelson WB, Slater S, Gold P & Gillin JC. The effect of growth hormone administration on human sleep: a dose-response study. *Biological Psychiatry* 1980 **15** 613–618.
- 143 Kern W, Halder R, al-Reda S, Späth-Schwalbe E, Fehm HL & Born J. Systemic growth hormone does not affect human sleep. *Journal of Clinical Endocrinology and Metabolism* 1993 **76** 1428–1432.
- 144 Astrom C. Interactions between sleep and growth hormone. Evaluated by manual polysomnography and automatic power spectrum analysis. *Acta Neurologica Scandinavica* 1995 **92** 281–296.
- 145 Wu RH & Thorpy MJ. Effect of growth hormone treatment on sleep EEGs in growth hormone deficient children. *Sleep* 1988 **11** 425–429.
- 146 Steiger A, Guldner J, Hemmeter U, Rothe B, Wiedemann K & Holsboer F. Effects of growth hormone-releasing hormone and somatostatin on sleep EEG and nocturnal hormone secretion in male controls. *Neuroendocrinology* 1992 **56** 566–573.
- 147 Kerkhofs M, Van Cauter E, Van Onderbergen A, Caufriez A, Thorner MO & Copinschi G. Sleep-promoting effects of growth hormone-releasing hormone in normal men. *American Journal of Physiology* 1993 **264** E594–E598.
- 148 Marshall L, Molle M, Boschen G, Steiger A, Fehm HL & Born J. Greater efficacy of episodic than continuous growth hormone releasing hormone (GHRH) administration in promoting slow wave sleep (SWS). *Journal of Clinical Endocrinology and Metabolism* 1996 **81** 1009–1013.
- 149 Guldner J, Schier T, Friess E, Colla M, Holsboer F & Steiger A. Reduced efficacy of growth hormone-releasing hormone in modulating sleep endocrine activity in the elderly. *Neurobiology of Aging* 1997 **18** 491–495.
- 150 Frieboes RM, Murck H, Schier T, Holsboer F & Steiger A. Somatostatin impairs sleep in elderly human subjects. *Neuropsychopharmacology* 1997 **16** 339–345.
- 151 Kupfer DJ, Jarrett DB & Ehlers CL. The effect of SRIF on the EEG sleep of normal men. *Psychoneuroendocrinology* 1992 **17** 37–43.
- 152 Obal F Jr, Floyd R, Kapas L, Bodosi B & Krueger JM. Effects of systemic GHRH on sleep in intact and hypophysectomized rats. *American Journal of Physiology* 1996 **270** E230–E237.
- 153 Weikel JC, Wichniak A, Ising M, Brunner H, Friess E, Held K *et al.* Ghrelin promotes slow-wave sleep in humans. *American Journal of Physiology – Endocrinology and Metabolism* 2003 **284** E407–E415.
- 154 Frieboes RM, Murck H, Maier P, Schier T, Holsboer F & Steiger A. Growth hormone-releasing peptide-6 stimulates sleep, growth hormone, ACTH and cortisol release in normal man. *Neuroendocrinology* 1995 **61** 584–589.
- 155 Frieboes RM, Murck H, Antonijevic IA & Steiger A. Effects of growth hormone-releasing peptide-6 on the nocturnal secretion of GH, ACTH and cortisol and on the sleep EEG in man: role of routes of administration. *Journal of Neuroendocrinology* 1999 **11** 473–478.
- 156 Copinschi G, Leproult R, Van Onderbergen A, Caufriez A, Cole KY, Schilling LM *et al.* Prolonged oral treatment with MK-677, a novel growth hormone secretagogue,

- improves sleep quality in man. *Neuroendocrinology* 1997 **66** 278–286.
- 157 Holsboer F, von Bardeleben U & Steiger A. Effects of intravenous corticotropin-releasing hormone upon sleep-related growth hormone surge and sleep EEG in man. *Neuroendocrinology* 1988 **48** 32–38.
- 158 Born J, Späth-Schwalbe E, Schwakenhofer H, Kern W & Fehm HL. Influences of corticotropin-releasing hormone, adrenocorticotropin, and cortisol on sleep in normal man. *Journal of Clinical Endocrinology and Metabolism* 1989 **68** 904–911.
- 159 Tsuchiyama Y, Uchimura N, Sakamoto T, Maeda H & Kotorii T. Effects of hCRH on sleep and body temperature rhythms. *Psychiatry and Clinical Neurosciences* 1995 **49** 299–304.
- 160 Born J, DeKloet ER, Wenz H, Kern W & Fehm HL. Gluco- and antimineralocorticoid effects on human sleep: a role of central corticosteroid receptors. *American Journal of Physiology* 1991 **260** E183–E188.
- 

Received 23 June 2003

Accepted 24 July 2003